

### IN THE CLAIMS

Please amend the claims as follows:

- 1-56. (Canceled).
57. (Currently Amended) A method for the stable transformation of monocot plant tissue or cells, comprising:
- a) selecting an amount of ~~a one or more~~ one or more ~~sulfhydryl-containing agent~~ sulfhydryl-containing agents effective in solid co-cultivation media to enhance the stable transformation of monocot plant tissue or cells with *Agrobacterium* relative to the stable transformation of monocot plant tissue or cells with *Agrobacterium* in the absence of the agent, wherein if the sulfhydryl-containing agent is cysteine, cysteine is present at a concentration of at least ~~[[50]]~~ 100 mg/L;
  - b) co-culturing on solid media monocot plant tissue or cells and an *Agrobacterium* containing a recombinant DNA, wherein the solid media comprises the amount of the one or more sulfhydryl-containing agents;
  - c) identifying stably transformed monocot plant tissue or cells; and
  - d) regenerating a transformed monocot plant from the stably transformed monocot plant tissue or cells.
58. (Original) The method of claim 57 or 62 wherein the efficiency of stable transformation in the presence of the agent is at least 10% greater than the efficiency of transformation in the absence of the agent.
59. (Previously Presented) The method of claim 57 or 62 wherein the efficiency of stable transformation in the presence of the agent is at least 0.5% greater than the efficiency of transformation in the absence of the agent.

- 
60. (Previously Presented) The method of claim 57 or 62 wherein the transformed tissue or cells are identified by selection.
61. (Original) The method of claim 60 wherein the transformed tissue or cells are selected for in hygromycin.
62. (Currently Amended) A method for the stable transformation of plant tissue or cells, comprising:
- a) selecting an amount of cysteine effective in solid co-cultivation media to enhance the stable transformation of plant tissue or cells with *Agrobacterium* relative to the stable transformation of plant tissue or cells with *Agrobacterium* in the absence of cysteine, wherein cysteine is present at a concentration of at least ~~[[50]]~~ 100 mg/L;
  - b) co-culturing on solid media plant tissue or cells and an *Agrobacterium* containing a recombinant DNA, wherein the solid media comprises the amount of cysteine;
  - c) identifying stably transformed plant tissue or cells; and
  - d) regenerating a transformed plant from the stably transformed plant tissue or cells.
63. (Original) The method of claim 57 or 62 wherein the stable transformation is enhanced by at least 5-fold.
64. (Original) The method of claim 57 or 62 wherein the stable transformation is enhanced by at least 10%.
- 65-66. (Canceled).
67. (Currently Amended) The method of claim 57 wherein ~~[[one]]~~ the agent is cysteine.
68. (Currently Amended) The method of claim 57 ~~[[or 62]]~~ wherein ~~[[one]]~~ the agent is glutathione, sodium thiosulfate, or dithiothreitol.

69-70. (Canceled).

71. (Original) The method of claim 57 or 62 wherein the recombinant DNA comprises a selectable marker.
72. (Original) The method of claim 57 or 62 wherein the recombinant DNA comprises a detectable marker.
73. (Original) The method of claim 57 or 62 wherein the recombinant DNA comprises a promoter operably linked to an open reading frame of interest.
74. (Original) The method of claim 68 wherein the glutathione is present at 0.4 g/L or 0.001 to 1 mM, sodium thiosulfate is present at 0.1 to 20 mM, or dithiothreitol is present at 1 g/L or 0.75 to 2 mM.
75. (Original) The method of claim 57 or 62 wherein the plant tissue or cells are maize, wheat, sugarcane or rice tissue or cells.
76. (Previous Presented) The method of claim 57 or 62 wherein the plant tissue or cells are maize, wheat or rice tissue or cells.
77. (Canceled)
78. (Currently Amended) The method of claim [[57 or]] 62 or 67 wherein cysteine is present at a concentration of at least 400 mg/L.